

# Limits on $\nu_\mu(\nu_e) \rightarrow \nu_\tau$ oscillations and tau neutrino magnetic moment from neutrino-electron scattering

S.N.Gninenko<sup>1</sup>

Institute for Nuclear Research, Russian Academy of Sciences,  
Moscow 117 312, Russia

## Abstract

The combined effect of neutrino flavor oscillations and neutrino magnetic moment on neutrino-electron scattering is discussed. It is shown, that if the tau neutrino has a (large) magnetic moment and can oscillate into a neutrino of another flavor it can be used for a sensitive search for  $\nu_\mu(\nu_e) \rightarrow \nu_\tau$  neutrino oscillations and the tau neutrino magnetic moment in neutrino -electron scattering experiments. The combined limits on the mixing angles and the tau neutrino magnetic moment  $\sin^2 2\theta_{\mu\tau} \times \mu_{\nu_\tau}^2 < 1.1 \times 10^{-18} \mu_B^2$  and  $\sin^2 2\theta_{e\tau} \times \mu_{\nu_\tau}^2 < 2.3 \times 10^{-18} \mu_B^2$  for  $\Delta m^2 > 10 \text{ eV}^2$  are presented based on results of a study of the neutrino-electron elastic scattering at LAMPF. For  $\nu_\mu \rightarrow \nu_\tau$  oscillations, this would results in sensitive limit of  $\sin^2 2\theta_{\mu\tau} < 4 \times 10^{-6}$ , assuming the tau neutrino magnetic moment being equal to the present experimental limit of  $5.4 \times 10^{-7} \mu_B$ . The tau neutrino magnetic moment would be constrained to  $\mu_{\nu_\tau} < 1.0 \times 10^{-8} \mu_B$ , assuming the existence of  $\nu_\mu \rightarrow \nu_\tau$  oscillations with the mixing angle  $\sin^2 2\theta_{\mu\tau}$  equal to the present experimental limit of 0.01 for  $\Delta m^2 > 10 \text{ eV}^2$ . Under similar assumptions the corresponding limits for  $\nu_e \rightarrow \nu_\tau$  oscillations could be set to  $\sin^2 2\theta_{e\tau} < 8 \times 10^{-6}$  and  $\mu_{\nu_\tau} < 3.9 \times 10^{-9} \mu_B$ .

In the Standard Model(SM) the properties of neutrino are strictly defined: it is a massless particle with zero electric charge and magnetic moment. Beyond the SM a nonzero mass of the neutrino is required for Grand Unified Theories (for review see e.g. [1]). In addition, massive tau neutrinos are natural candidates for the hot component of the dark matter of the Universe. It is generally assumed that massive neutrinos could manifest themselves through the effect of neutrino oscillations first postulated by Pontecorvo [2] a long time ago. Nonzero electromagnetic properties of the neutrino have also been discussed in many extensions of the SM [1]. In its simplest extension, for example, neutrinos may have masses and acquire a magnetic moment through radiative corrections [3, 4]:

$$\mu_{\nu_i} = \frac{3eG_F m_{\nu_i}}{8\sqrt{2}\pi^2} = 3.2 \times 10^{-19} (m_{\nu_i}/1 \text{ eV}) \mu_B \quad (1)$$

---

<sup>1</sup>E-mail address: Sergei.Gninenko@cern.ch

where  $G_F$  is the Fermi constant,  $\mu_{\nu_i}$  ( $i = 1, 2, 3$ ) is the neutrino magnetic moment,  $m_{\nu_i}$  is the Dirac neutrino mass eigenstate and  $\mu_B = e/2m_e$  is the Bohr magneton. This model, however predicts neutrino magnetic moments which are too small to give any observable effects discussed in the present paper. Some models predict for the neutrino magnetic moment values as large as  $10^{-10}\mu_B$ , see e.g. ref.[5], which are large enough to be observed through neutrino electromagnetic interactions.

At present the  $\nu_\mu \rightarrow \nu_\tau$  and  $\nu_e \rightarrow \nu_\tau$  neutrino oscillations are a subject of a search at the CERN SPS wide-band neutrino beam by the CHORUS and NOMAD experiments [6, 7]. These searches are based on the appearance of tau neutrinos in a beam of muon neutrinos and requires detection of the  $\nu_\tau$  charged current interaction

$$\nu_\tau N \rightarrow \tau^- X \quad (2)$$

The intrinsic fraction of  $\nu_\tau$ 's in the beam is expected to be negligibly small ( $\sim 10^{-6}$ ). Two methods are used to detect reaction (2). The CHORUS collaboration adopted the method based on the direct observation the  $\tau$  decay topology. In the NOMAD experiment a search of an apparent non-conservation of transverse momentum due to undetected neutrinos in  $\tau$  decays is used [8].

In this note we show that combined existence of  $\nu_\mu \rightarrow \nu_\tau$  and/or  $\nu_e \rightarrow \nu_\tau$  oscillations and a (large) nonzero magnetic moment of the tau neutrino would increase the total rate of events in  $\nu_\mu(\nu_e)$  neutrino- electron scattering experiments. This effect could be used for a sensitive search for these combined properties of the tau neutrino in the reactions of neutrino - electron scattering.

Assuming that a muon neutrino beam has a component of tau neutrinos due to  $\nu_\mu \rightarrow \nu_\tau$  oscillations. Then if magnetic moment of the  $\nu_\tau$  exists, it will contribute to a non-coherent part of the  $\nu_\tau e^-$  scattering cross section via the reaction that change the helicity of the tau neutrino (hence right-handed neutrino states should exist) [9]. This might result in observable deviations from purely  $\nu_\mu e^-$  electro-weak reaction which is well predicted by the SM. Indeed, since the electromagnetic cross section is orders of magnitude larger than the weak cross section, even a small fraction of tau neutrinos with nonzero magnetic moment in the muon neutrino beam could lead to an observable effects in  $\nu_\mu e^-$  scattering, while the magnetic moment of muon neutrino could be small enough to contribute effectively to  $\nu_\mu e^-$  scattering.

The neutrino-electron scattering process via magnetic moment has a cross section [9]:

$$\frac{d\sigma_\mu}{dy} = \frac{\pi\alpha^2}{m_e^2} \frac{\mu_\nu^2}{\mu_B^2} \frac{1-y}{y} \quad (3)$$

where  $y = E_e/E_\nu$ , and  $\mu_\nu$  is the neutrino magnetic moment. The production rate of isolated electrons via  $\nu_\tau e^-$  scattering in the detector depends also on the probability to find a  $\nu_\tau$  neutrino in the neutrino beam. This probability can be calculated from the neutrino survival and transition probabilities.

Consider first the case of a mixture of three massive neutrinos

$$\nu_l = \sum_{i=1}^3 U_{li} \nu_i, \quad (4)$$

Here  $\nu_l, \nu_i$  are the fields of the neutrino with flavor  $l$  and mass  $m_i$ , respectively and  $U$  is a unitary mixing matrix. Ignoring possible  $CP$  violation, for the probability of the transition  $\nu_l \rightarrow \nu_{l'}$  we have[10]

$$P(\nu_l \rightarrow \nu_{l'}) = \left| \delta_{ll'} - 4 \sum_{i>j} U_{l'i} U_{l'j} U_{li} U_{lj} \times \sin^2 \frac{\Delta m_{ij}^2 L}{4E_\nu} \right| \quad (5)$$

where  $l, l' = e, \mu, \tau$ ,  $L$  is the distance between the detector and the neutrino source and  $\Delta m_{ij}^2 = |m_i^2 - m_j^2|$ .

In case of two-neutrino mixing, e.g.  $\nu_\mu \rightarrow \nu_\tau$ , neutrino states evolve with a time  $t$  as

$$|\nu > (t) = a(t)|\nu_\mu > + b(t)|\nu_\tau > \quad (6)$$

where  $|\nu_\mu >$  and  $|\nu_\tau >$  denote weak eigenstates of  $\nu_\mu$  and  $\nu_\tau$  neutrinos, and  $a^2(t)$ ,  $b^2(t)$  are the probabilities to find  $\nu_\mu$  or  $\nu_\tau$  in the beam at a given moment  $t$ , respectively. It is assumed that  $a^2(0) = 1$  at  $t = 0$ . The probability  $b^2(t)$  depends on the parameters of  $\nu_\mu - \nu_\tau$  oscillations as :

$$b^2(t \simeq \frac{L}{c}) = P(\nu_\mu \rightarrow \nu_\tau) = \sin^2 2\theta_{\mu\tau} \sin^2 \frac{\Delta m^2 L}{4E} \quad (7)$$

or

$$P(\nu_\mu \rightarrow \nu_\tau) \approx \sin^2 2\theta_{\mu\tau} \sin^2 \frac{1.27 \Delta m^2 (eV^2) L (km)}{E (GeV)} \quad (8)$$

where  $\sin^2 2\theta_{\mu\tau}$  is the mixing angle, and  $\Delta m^2 = \Delta m_{32}^2$ .

Now, if an experiment sets a limit on a magnetic moment of the neutrino  $\mu_\nu < \mu_0$ , we may write in the general form the following equation, taking into account neutrino oscillations between different neutrino flavors and nonzero magnetic moment of the neutrinos:

$$\sum_{l, l' = e, \mu, \tau} f_{\nu_l} \times [1 - P(\nu_l \rightarrow \nu_{l'})] \times \mu_{\nu_l}^2 + f_{\nu_l} \times P(\nu_l \rightarrow \nu_{l'}) \times \mu_{\nu_{l'}}^2 \leq \mu_0^2 \quad (9)$$

where  $f_{\nu_l}$  is a factor corresponding to the flux of the  $\nu_l$ -neutrino at the source, and the fluxes are normalized to  $1 = \sum f_{\nu_l}$ . Here we assume that detection efficiencies for electrons produced in  $\nu_l e^-$  scattering are the same for neutrino of all three flavors. Note, that limit (9) is set based on an agreement between observed cross section in  $\nu e^-$  scattering experiment and SM expectations. The  $\nu_\mu \rightarrow \nu_e$  oscillations could also affect the observed  $\nu e$  cross section because of the difference in cross sections for  $\nu_e e^-$  and  $\nu_\mu (\nu_\tau) e^-$  scattering. However, according to the present data on  $\nu_\mu \rightarrow \nu_e$  oscillation from the LSND experiment,  $P(\nu_\mu \rightarrow \nu_e) \approx 3 \times 10^{-3}$  for (large)  $\Delta m^2 > 10 \text{ eV}^2$  [11]. This

could shift the  $\nu_\mu e^-$  cross section by at most 1%, which is small in comparison with the statistical and systematic errors of the experiments discussed below. More detailed discussions of the effect of neutrino flavor oscillations on  $\nu e^-$  scattering can be found in ref. [12]. We will ignore this effect to avoid complication of our formula (9).

The best limit on neutrino magnetic moment was obtained from the experiment on study of the  $\nu_e e^-$  elastic scattering at LAMPF, ref. [13, 14], and was confirmed by the CHARM II experiment at CERN at much higher energy, ref.[15]. The average  $L/E$  ratio was  $\approx 0.3$  in ref.[13, 14], and  $\approx 0.05$  in ref.[15].

The upper limits on  $\mu_{\nu_\mu}$  and  $\mu_{\nu_e}$  from the LAMPF experiment was obtained using the following equation ( ref.[14], Eq.(23))

$$\mu_{\nu_e}^2 + 2.1 \times \mu_{\nu_\mu}^2 \leq \mu_0^2 = 1.16 \times 10^{-18} \mu_B^2 \quad (10)$$

and therefore

$$\mu_{\nu_\mu} < 7.4 \times 10^{-10} \mu_B \quad (\mu_{\nu_e} = 0) \quad (11)$$

$$\mu_{\nu_e} < 10.8 \times 10^{-10} \mu_B \quad (\mu_{\nu_\mu} = 0) \quad (12)$$

Taking into account  $\nu_\mu(\nu_e) \rightarrow \nu_\tau$  oscillations Eq.(10) can be rewritten in the form similar to Eq.(9)

$$\begin{aligned} & [1 - P(\nu_e \rightarrow \nu_\tau)] \times \mu_{\nu_e}^2 + P(\nu_e \rightarrow \nu_\tau) \times \mu_{\nu_\tau}^2 \\ & + 2.1 \times [1 - P(\nu_\mu \rightarrow \nu_\tau)] \times \mu_{\nu_\mu}^2 + 2.1 \times P(\nu_\mu \rightarrow \nu_\tau) \times \mu_{\nu_\tau}^2 \leq \mu_0^2 \end{aligned} \quad (13)$$

which can be used to obtain combined limits on the probability of  $\nu_\mu(\nu_e) \rightarrow \nu_\tau$  oscillations versus tau neutrino magnetic moment:

$$P(\nu_e \rightarrow \nu_\tau) \times \mu_{\nu_\tau}^2 + 2.1 \times P(\nu_\mu \rightarrow \nu_\tau) \times \mu_{\nu_\tau}^2 \leq 1.16 \times 10^{-18} \mu_B^2 \quad (\mu_{\nu_e} = \mu_{\nu_\mu} = 0) \quad (14)$$

$$P(\nu_e \rightarrow \nu_\tau) \times \mu_{\nu_\tau}^2 \leq 1.16 \times 10^{-18} \mu_B^2 \quad (P(\nu_\mu \rightarrow \nu_\tau) = 0, \mu_{\nu_e} = \mu_{\nu_\mu} = 0) \quad (15)$$

$$P(\nu_\mu \rightarrow \nu_\tau) \times \mu_{\nu_\tau}^2 \leq 5.5 \times 10^{-19} \mu_B^2 \quad (P(\nu_e \rightarrow \nu_\tau) = 0, \mu_{\nu_e} = \mu_{\nu_\mu} = 0) \quad (16)$$

From Eqs.(15,16) combined limits on the mixing angles and tau neutrino magnetic moment for  $\Delta m^2 > 10 \text{ eV}^2$ , which corresponds to the large  $\Delta m^2$  region in the LAMPF experiment can be derived

$$\sin^2 2\theta_{e\tau} \times \mu_{\nu_\tau}^2 \leq 2.3 \times 10^{-18} \mu_B^2 \quad (17)$$

$$\sin^2 2\theta_{\mu\tau} \times \mu_{\nu_\tau}^2 \leq 1.1 \times 10^{-18} \mu_B^2 \quad (18)$$

Assuming the tau-neutrino magnetic moment to be equal to the present experimental limit of  $5.4 \times 10^{-7} \mu_B$  from the BEBC experiment at CERN, ref.[16], the following limits

on the oscillation probabilities  $P(\nu_e \rightarrow \nu_\tau)$ ,  $P(\nu_\mu \rightarrow \nu_\tau)$  and mixing angles  $\sin^2 2\theta_{e\tau}$ ,  $\sin^2 2\theta_{\mu\tau}$  can be obtained from Eqs.(15,16,17,18), respectively

$$P(\nu_e \rightarrow \nu_\tau) < 4 \times 10^{-6} \quad (19)$$

$$\sin^2 2\theta_{e\tau} < 8 \times 10^{-6} \quad (20)$$

$$P(\nu_\mu \rightarrow \nu_\tau) < 2 \times 10^{-6} \quad (21)$$

$$\sin^2 2\theta_{\mu\tau} < 4 \times 10^{-6} \quad (22)$$

Under assumption of the existence of neutrino oscillations with mixing angles equal to the present experimental limits of  $\sin^2 2\theta_{e\tau} = 0.15$  obtained from  $\overline{\nu}_e \rightarrow \nu_x$  disappearance reactor experiments in Bugey [17] and in Krasnoyarsk [18], or  $\sin^2 2\theta_{\mu\tau} = 0.01$  for  $\nu_\mu \rightarrow \nu_\tau$ , obtained from the exclusion plot of ref.[19] for  $\Delta m^2 > 10 \text{ eV}^2$ , the following upper bounds on the magnetic moment of the tau neutrino can be derived from Eqs.(17,18):

$$\mu_{\nu_\tau} < 3.9 \times 10^{-9} \mu_B \quad (\nu_e \rightarrow \nu_\tau) \quad (23)$$

$$\mu_{\nu_\tau} < 1.0 \times 10^{-8} \mu_B \quad (\nu_\mu \rightarrow \nu_\tau) \quad (24)$$

In (23), no distinction has been made between  $\nu_e$  and  $\overline{\nu}_e$ . The exclusion regions in the  $(\sin^2 2\theta_{e\tau}, \mu_{\nu_\tau}/\mu_B)$  and  $(\sin^2 2\theta_{\mu\tau}, \mu_{\nu_\tau}/\mu_B)$  planes obtained from Eqs.(17,18) are shown in Figure 1,2, respectively. Experimentally excluded regions for  $\mu_{\nu_\tau}$ ,  $\sin^2 2\theta_{e\tau}$  and  $\sin^2 2\theta_{\mu\tau}$  are also indicated.

Assuming the existence of  $\nu_\mu \rightarrow \nu_\tau$  neutrino oscillations with mixing angles  $\sin^2 2\theta_{\mu\tau}$  equal to the present experimental limits of  $3.5 \times 10^{-3}$  from the CHORUS [20], or  $3.7 \times 10^{-3}$  from the NOMAD [21] experiments, the limit of

$$\mu_{\nu_\tau} < 1.8 \times 10^{-8} \mu_B \quad (\nu_\mu \rightarrow \nu_\tau) \quad (25)$$

can also be obtained, however for the mass region  $\Delta m^2 > 1000 \text{ eV}^2$ .

Using the upper limit on muon neutrino magnetic moment from the CHARM II experiment [15] one can obtain limits on  $P(\nu_\mu \rightarrow \nu_\tau)$ ,  $\sin^2 2\theta_{\mu\tau}$  and  $\mu_{\nu_\tau}$  of the same order as the limits (21),(22), and (24), respectively. Note that to separate an oscillation signal from magnetic moment scattering in  $\nu e^-$  reaction one can look at its energy or distance dependence. For example, the probability of oscillations changes as a  $E_\nu^{-2}$  for small  $\Delta m^2$ , while the integral cross section for the magnetic moment scattering varies only logarithmically with  $E_\nu$ .

## Acknowledgements

The author wishes to thank L. Camilleri, L. DiLella, E. do Couto e Silva, V.A. Kuzmin and A. Rubbia for useful discussions and comments. Communications with K. Winter on latest CHORUS results are gratefully acknowledged.

## References

- [1] R.N. Mohapatra and P.B. Pal, in “Massive Neutrinos in Physics and Astrophysics”, World Scientific, Singapore, 1991.
- [2] B. Pontecorvo, Zh. Eksp. Teor. Fiz. **33** (1957) 549; [JETP **6**(1958)429].
- [3] B.W. Lee and R.E. Shrock, Phys. Rev. **D16** (1977) 1444.
- [4] W. Marciano and A.I. Sanda, Phys. Lett. **B67** (1977) 303.
- [5] M.B. Voloshin, M.I. Vysotskii and L.B. Okun, Zh. Eksp. Teor. Fiz. **91**(1986) 754.
- [6] E. Eskut et al., CHORUS Collaboration, Nucl. Instr. and Meth. **A401**(1997)7.
- [7] J. Altegoer et al., NOMAD Collaboration, to appear in Nucl. Instr. and Meth.
- [8] C. Albright and E. Shrock, Phys. Lett. **B84** (1979) 123.
- [9] G. Domogatskii and D. Nadezhin, Sov. J. Nucl. Phys. **12**(1971)678.
- [10] See e.g. review: S.M. Bilenky and B. Pontecorvo, Phys. Rep. **41** (1978) 225.
- [11] C. Athanassopoulos et al., Phys. Rev. **C54** (1996) 2685;  
C. Athanassopoulos et al., UCRHEP-E191/LA-UR-97-1998, submitted to Phys. Rev. C.
- [12] W.J. Marciano, Phys. Rev. **D36** (1987) 2859.
- [13] D.A. Krakauer et al., Phys. Lett. **B252** (1990) 177.
- [14] R.C. Allen et al., Phys. Rev. **D47** (1993) 11.
- [15] P. Vilain et al., Phys. Lett. **B345** (1995) 115.
- [16] A.M. Cooper - Sarkar et al., Phys. Lett. **B280** (1992) 153.
- [17] B. Achkar et al., Nucl. Phys. **434** (1995) 503.
- [18] G.S. Vidyakin et al., JETP Lett. **59** (1994) 390.
- [19] N. Ushida et al., Phys. Rev. Lett. **57** (1986) 2897.
- [20] CHORUS collaboration, CERN-PPE/97-149, submitted to Phys. Lett. B (1998);  
The latest limit of  $\sin^2 2\theta_{\mu\tau} < 2.3 \times 10^{-3}$  (90% *C.L.*) for large  $\Delta m^2$  was recently reported: J. Brunner, Talk at CERN Seminar, October, 1997.
- [21] A. Rubbia, “Neutrino Physics at Accelerators”, Talk given at Lepton-Photon Symposium, Hamburg, 1997.




Figure 1: The exclusion regions for combination of  $\nu_e \rightarrow \nu_\tau$  oscillations and tau neutrino magnetic moment  $\mu_{\nu_\tau}$  in the  $(\sin^2 2\theta_{e\tau}, \mu_{\nu_\tau}/\mu_B)$  plane. Light shaded area is excluded by this analysis based on the LAMPF results on  $\nu e^-$  scattering [13, 14]. Dark shaded area corresponds to the experimentally excluded regions  $\mu_{\nu_\tau} > 5.4 \times 10^{-7} \mu_B$  [16] and  $\sin^2 2\theta_{e\tau} > 0.15$  [17, 18]. The limits are valid for  $\Delta m^2 > 10 \text{ eV}^2$ .




Figure 2: The exclusion regions for combination of  $\nu_\mu \rightarrow \nu_\tau$  oscillations and tau neutrino magnetic moment  $\mu_{\nu_\tau}$  in the  $(\sin^2 2\theta_{\mu\tau}, \mu_{\nu_\tau}/\mu_B)$  plane. Light shaded area is excluded by this analysis based on the LAMPF results on  $\nu e^-$  scattering [13, 14]. Dark shaded area corresponds to the experimentally excluded regions  $\mu_{\nu_\tau} > 5.4 \times 10^{-7} \mu_B$  [16] and  $\sin^2 2\theta_{\mu\tau} > 0.01$  [19]. The limits are valid for  $\Delta m^2 > 10 \text{ eV}^2$ .